

The jumping spiders (Araneae, Salticidae) of Khabarovsk Province (Russian Far East)

Dmitrii V. Logunov & Wanda Wesołowska

Logunov, D. V., Zoological Museum, Institute of Biology, the Russian Academy of Sciences, Frunze street 11, 630091 Novosibirsk, Russia

Wesołowska, W., Zoological Institute, Wrocław University, Sienkiewicza 21, 50-335 Wrocław, Poland

Received 19 February 1992, accepted 18 June 1992

The paper presents data on the Salticidae (except of the genus *Euophrys*) of Khabarovsk Province (Russian Far East). Thirty-six species are reported; among them three are described as new (*Harmochirus nigriculus*, *Marpissa dersuzalai*, *Marpissa zebra*), four are new for Russia (*Harmochirus pullus*, *Marpissa pulchra*, *Phintella linea*, *P. parva*) and twelve are new for Khabarovsk Province (*Harmochirus latens*, *Evarcha albaria*, *Heliophanus dubius*, *Marpissa dybowski*, *M. elongata*, *M. pulla*, *Myrmarachne formicaria*, *M. lugubris*, *Phintella arenicolor*, *P. popovi*, *Salticus cingulatus*, *Sitticus cutleri*). Two names are synonymized (*Yaginumaella ususudi* with *Y. striatipes* and *Phintella mellotei* with *P. arenicolor*). The neotype of *Phintella castrisiana* is designated. *Bianor latens* is assigned to the genus *Harmochirus*, and the first description of the male of this species is provided. The differentiating diagnoses of poorly known Eastern Palaeartic species of the genera *Marpissa*, *Myrmarachne* and *Phintella* are given.

1. Introduction

Our knowledge of the Salticidae from different parts of the Palaeartic is very biased; whereas the Western Palaeartic species are relatively well known, our knowledge of the Central and Eastern Palaeartic species is grossly inadequate.

The current distribution of the Palaeartic species was shaped to a large extent by the Ice Age. After the retreat of the glaciers the species recolonized the Palaeartic from the refugia. Prószyński (1976) put forward a hypothesis that

the species spreading from Western Palaeartic refugia were more expansive, colonizing large areas, and reaching in some cases to the shores of the Pacific Ocean. On the other hand, the species from the Eastern centres of dispersal were less dynamic, and this resulted in the small ranges of the Eastern Palaeartic species. To verify this hypothesis one needs to have a much better idea of species distribution patterns in the Central and Eastern Palaeartic. This demands not only faunistic studies in poorly known areas, but taxonomic work as well. The Asiatic species are




Fig. 1. Map of study area. — 1: Khabarovskii Krai. — 1a: Evreiskaya Autonoma Oblast (separate administrative unit constituting part of Khabarovskii Krai). — 2: Amurskaya Oblast. — 3: Primore.

insufficiently known, and for many of them good descriptions and diagnoses are lacking, making their correct identification practically impossible.

Our knowledge of the Salticidae of the Khabarovsk Province (Khabarovskii Krai) (Fig. 1) stems from only a few publications. Initial information was provided by Grube (1861), who reported eight species from there. It was not until more than a century later that papers by Prószyński (1979) and Dunin (1984) allowed us to extend the list of species known from this area to twenty. Four of the species from this list have not been found in our material. These are: *Evarcha laetabunda* (C. L. Koch), *Pellenes ignifrons* (Grube), *Sitticus caricis* (Westring) and *Sitticus pubescens* (Fabricius). The latter species was recorded in the Amur river valley by Grube (1861). This information was repeated by Kharitonov (1932), Azheganova & Stenchenko (1977), Dunin (1984) and Nenilin (1985). Unfortunately, the specimen has disappeared from Grube's collection (Prószyński 1971), so verification of this record is impos-

ible. As this European species has never again been found East of the Urals (see Prószyński 1983: fig. 5), it seems that its inclusion in the fauna of Khabarovskii Krai is based on a mistake.

This paper is based on the rather large amount of material collected by D. V. Logunov in Bolshekhkhechtsirskii reserve (referred to as BR in the text), situated 15–20 km South of Khabarovsk, in 1987. Smaller collections were accumulated by D. K. Kurenshchikov and Dr. N. A. Ryabinin (Institute of Water and Ecological Problems, Khabarovsk) and of S. V. Ivanov (Bolshekhkhechtsirskii reserve). The material in the Zoological Museum of Moscow University (collection of V. I. Sychevskaya) and in the Museum of Natural History in Wrocław (collection of A. E. Grube) was studied as well.

Both the previous information and our data refer to the southern part of Khabarovskii Krai, mainly the Amur river valley (see Fig. 1). For some species data from the neighbouring Amurskaya Oblast and Primore are included as well. For *Phintella popovi* and *Sitticus cutleri* data from Khakassya (South of Krasnoyarskii Krai) and for *Harmochirus latens* data from Khakassya and Buryatya are added too.

All the Salticidae species apart from those of the genus *Euophrys* are dealt with here. The latter genus will be treated separately, as its Eastern Palaearctic species demand a thorough taxonomic revision. Most probably 2–3 *Euophrys* species occur within the study area.

The material forming the foundations of this paper originates from the following collections:

- | | |
|------|---|
| BI | Zoological Museum of Institute of Biology, the Russian Academy of Sciences, Novosibirsk |
| MNH | Museum of Natural History, Wrocław University, Wrocław |
| ZMMU | Zoological Museum of Moscow University, Moscow |
| ZI | Zoological Institute, the Russian Academy of Sciences, St. Petersburg |
| IWP | Institute of Water and Ecological Problems, the Russian Academy of Sciences, Khabarovsk |
| IBS | Biological Institute of WSRP, Siedlce |
| ZMT | Zoological Museum, Turku University, Turku. |

2. Methods

Specimens were examined in a dish with alcohol. Descriptions of colours pertain to wet specimens. The drawings were made with the aid of a reticular eyepiece attached to a stereomicroscope with a magnification of 7.5 to 48 times. If not stated otherwise, specimens from Khabarovskii Krai are shown in the figures. The male pedipalp and epigyne were removed for the study. The epigyne was cleared in warm lactic acid or macerated in 10% KOH for 48 hours at normal room temperature and cleared in xylene. After drawing, the genitalia were placed in microvials with alcohol.

All measurements are given in mm. In the descriptions the following abbreviations are used:

- AM diameter of anterior median eyes
 AL distance between anterior lateral eyes
 PL distance between posterior eyes.

3. Taxonomic survey of species

Aelurillus festivus (C. L. Koch, 1834)

Material: Khabarovskii Krai, BR, the Ussuri river, 2♂, 1♀, 10.IX.1987, S. Ivanov (BI, No 150, 155).

Habitat. Collected on rocky river shore among stones.

Distribution. Widely distributed Palaearctic species, recorded from the Russian Far East by Grube (1861), Kulczyński (1895), Prószyński (1976: map 1, 1979), Dunin (1984), Nenilin (1985) and Shternbergs (1988).

Carrhotus xanthogramma (Latreille, 1819)

Material: Khabarovskii Krai, BR, 2♂, 5♀, 5–19.VI. 1987, D. Logunov (BI, No 690, 694, 898, 900), 1♂, 1♀ (MNH), 1♀, 8–10.VIII.1987, S. Ivanov (BI); BR, Bychikha village, 2♂, 3♀, 2–31.V.1988, S. Ivanov (BI, No 692), 2♂ (MNH); Nanaiskii distr., Slavyanka, 1♀, 1983, N. Ryabinin (ZMMU); the Amur river, 2♀, [leg.] Maack, coll. Grube (MNH).

Habitat. This species was collected by sweeping grass in deciduous woods (aspen, birch, oak); also recorded in coastal vegetation on stream submersion areas.

Distribution. Widely distributed Palaearctic species, recorded from the Russian Far East by Kulczyński (1895), Prószyński (1971, 1976: map 66, 1979), Dunin (1984), Nenilin (1985) and Shternbergs (1988).

Dendryphantus fusconotatus (Grube, 1861) ?

Material: Khabarovskii Krai, BR, 2♂, 17–19.VI.1987, D. Logunov (BI, No 1117, 1118).

Distribution. Eastern Palaearctic species, recorded repeatedly in Khabarovskii Krai and Primore (Grube 1861, Kharitonov 1932, Prószyński 1971, 1976: map 44, 1979, Azheganova & Stenchenko 1977, Izmailova 1980, Dunin 1984, Nenilin 1985 and Marusik 1988).

These specimens differ slightly from the type. Perhaps they belong to another, closely related species. Settlement of this question will be possible only after thorough study.

Evarcha albaria (L. Koch, 1878)

Material: Khabarovskii Krai, BR, 7♂, 1♀, 2–19.VI. 1987, D. Logunov (BI, No 678, 681), 2♂, 2♀ (MNH), 1♂, 8–10.VIII.1987, S. Ivanov (BI, No 680).

Habitat. Collected by sweeping grass and bushes in deciduous forest (aspen, birch, oak), as well as in litter.

Distribution. Eastern Palaearctic species. New for Khabarovskii Krai, previously recorded in the Russian Far East only from Primore (Kulczyński 1895, Kharitonov 1932, Prószyński 1976: map 128, 1979, Dunin 1984, Nenilin 1985).

Evarcha arcuata (Clerck, 1757)

Material: Khabarovskii Krai, BR, 14♂, 6♀, 2–19.VI. 1987, D. Logunov (BI, No 280, 291, 316, 321, 323, 324, 327), 1♂, 1♀ (MNH), 1♀ (IBS); the Amur river, Rybachii island, 2♀, 21.VI.1987, D. Kurenschikov (BI, No 317).

Habitat. Collected by sweeping both in dry deciduous woods (aspen, birch, oak) and in moister places (among sparse ferns in alder carr with single larches), as well as in clearings and on meadows.




Fig. 2. *Harmochirus latens* (Logunov), male. — A–B: palpal organ, ventral and lateral views. — C: cheliceral dentition. — D: first leg.

Distribution. Widely distributed Palearctic species, recorded from the Russian Far East by Prószyński (1976: map 130, 1979), Azheganova & Stenchenko (1977), Dunin (1984), Nenilin (1985) and Marusik (1988).

***Evarcha falcata* (Clerck, 1757)**

Material: Khabarovskii Krai, BR, 2♂, 6♀, 15–22.VI.1987, D. Logunov (BI, No 181, 197, 201, 202, 205).

Habitat. Collected by sweeping in deciduous woods (aspen, birch, oak) and at wood edges.

Distribution. Widely distributed Holarctic species, recorded repeatedly in the Russian Far East (Kulczyński 1895, Prószyński 1976: map 131, 1979, Azheganova & Stentchenko 1977, Dunin 1984, Nenilin 1985).

***Harmochirus latens* (Logunov, 1991) comb. n.**
Figs. 2–3

Bianor latens Logunov, 1991

Material: Khabarovskii Krai, BR, 2♂, 10–19.VI.1987, D. Logunov (BI, No 645, 646); Amurskaya Oblast, Khin-

ganskii reserve, Antonovskoe forest-range, 1♀, 1.VIII.1983, Yu. Marusik (ZI); Primore, Khasanskii distr., near Primorskaya station, 1♂, 1♀, 31.V.1978, B. Zakharov (ZI); Buryatya, Selenginskii distr., Tazhnoe village, 1♀, 13.VIII.1984, B. Zakharov (BI, No 644); South of Krasnoyarskii Krai, Khakassya, Altaiskii distr., 40 km SE of Belyi Yar, 15–17 km E of Novorossiiskoe, the Jenisej river, 1♂, 24.VI.1990, D. Logunov (BI, No 1115).

Diagnosis. The shape of epigyne and swollen tibiae of front legs in males suggest that this species should be placed in the genus *Harmochirus*. Exact delimitation of *Bianor* and *Harmochirus*, though, calls for future detailed studies. The male similar to *Bianor aurocinctus* (Ohlert) and *Bianor aemulus* (Gertsch), may be distinguished by stronger tibial apophysis, flatter tegulum and position of seminal ducts (Fig. 2A–B). The female may be easily recognized by long sideward depressions of epigyne (Fig. 3A–B).

Description. Measurements (male/female). Cephalothorax: length 1.47/1.89, width 1.10/1.40, height 0.61/0.77. Abdomen: length 1.50/2.57, width 1.07/1.76. Eye field: length 0.83/1.03, AL 0.89/1.13, PL 1.07/1.46. AM 0.29/0.37. Legs: I 0.79/0.97 + 0.51/0.56 + 0.54/0.63 + 0.40/0.47 + 0.34/0.37; II 0.57/0.81 + 0.37/0.50 + 0.30/0.44 +




Fig. 3. *Harmochirus latens* (Logunov), female, epigyne and its internal structures. — A, C: specimen from Primore. — B, D: specimen from Amurskaya Oblast.

0.29/0.40 + 0.29/0.34; III 0.73/1.00 + 0.46/0.49 + 0.34/0.46 + 0.36/0.53 + 0.29/0.34; IV 0.71/1.03 + 0.36/0.50 + 0.41/0.60 + 0.43/0.63 + 0.29/0.37. — Male. Carapace flattened, much swollen in the area of eyes III, dark brown, eye field with metallic shine. Scarce brown bristles on carapace, few white scales behind posterior eyes. Clypeus low with long hairs. Chelicerae (Fig. 2C), maxillae, labium and sternum brown. Abdomen rounded, dark brown with gleaming scutum. Legs I thick dark brown with long dense black hairs on ventral surface of femora and tibiae (Fig. 2D). Remaining legs pale brown. Pedipalp dark with single straight tibial apophysis, bulbus more or less rounded, embolus very thin (Fig. 2A–B). — Female. Carapace dark brown, eye field black. Sides on cephalothorax and clypeus covered with white hairs. Sternum brown

with white hairs. Labium, maxillae and chelicerae brown. Abdomen and spinnerets dark grey. Legs I with thickened dark brown femora, patellae reddish-brown or brown, tibiae red with brown band on their distal parts, metatarsi and tarsi yellow or yellowish-grey. Remaining legs brown, their distal parts yellow or yellowish-grey. Pedipalp yellow or grey, its femur brown. Epigyne is given in Fig. 3A–B, its internal structures in Fig. 3C–D.

Habitat. Collected by sweeping grass in woods and floodplains, as well as in litter in poplar woods.

Distribution. Siberia: Khakassya, Tuva, Buryatya, Khabarovskii Krai, Amurskaya Oblast, Primore.

This species was described by Logunov (1991) on the basis of a female from Tuva. The male is described here for the first time.




Fig. 4. *Harmochirus nigriculus* sp. n. and *Bianor aurocinctus* (Ohlert), males. — A–B: *H. nigriculus*, holotype, palpal organ, ventral and lateral views. — C–D: *B. aurocinctus*, specimen from Sakhalin, palpal organ, ventral and lateral views. — VK: ventral knob on palpal tibia. — PD: pear-shaped depression in the tegulum cone.

***Harmochirus nigriculus* sp. n.**

Figs. 4–5

Material: Holotype: Khabarovskii Krai, environs of Khabarovsk, the Amur river, Zelonyi island, 1♂, 29.VI.1936, V. Sychevskaya (ZMMU, Ta-4658). — Paratypes: Khabarovskii Krai, Nanaiskii distr., the Amur river, Slavyanskii island, 1♀, 20.VI.1989, D. Kurenshchikov (BI, No 647); Amurskaya Oblast, Tambovskii distr., Kozmodemyanovka village, 3♀, VI.1990, G. Ganin (BI, No 1116); Primore, Skotovskii distr., Kangauz, 1♀ (incorrectly determined as “*B. aurocinctus*”), 12.VII.1968, V. Popov (ZI), 1♀ (incorrectly determined as “*B. aurocinctus*”), IX.1968, N. Ustimenko (ZI).

Diagnosis. The species clearly shows affinities with *Bianor aurocinctus* (Ohlert). The male may be distinguished by more clearly marked pear-shaped depression in tegulum cone (PD, cf. Fig. 4B and 4D), by position of turn of seminal duct, presence and situation of ventral knob on palpal tibia (VK, cf. Fig. 4A and 4C), and by smaller size of both pedipalp and whole body. The female can be separated by position of loops of insemination ducts (cf. Fig. 5B and 5C). It seems also to have differently coloured legs: *H. nigriculus* has brown patellae I and greyish-brown




Fig. 5. *Harmochirus nigriculus* sp. n. and *Bianor aurocinctus* (Ohlert), females. — A–B: *H. nigriculus*, paratype, epigyne and its internal structures. — C: *B. aurocinctus*, specimen from Sakhalin, internal structures of epigyne.

legs II–IV, while *B. aurocinctus* has red patellae I and yellow legs II–IV (see also Logunov & Marusik 1991).

Description. Measurements (male/female). Cephalothorax: length 1.42/1.73, width 1.10/1.28, height 0.56/0.75. Abdomen: length 1.71/2.50, width 1.19/1.50. Eye field: length 0.86/0.95, AL 0.91/1.05, PL 1.10/1.30. AM 0.29/0.34. Legs: I 1.09/0.93 + 0.64/0.58 + 0.68/0.60 + 0.64/0.35 + 0.34/0.35; II 0.61/0.70 + 0.36/0.45 + 0.38/0.40 + 0.34/0.40 + 0.48/0.28; III 0.79/0.93 + 0.37/0.50 + 0.34/0.43 + 0.57/0.45 + 0.29/0.30; IV 0.83/0.93 + 0.37/0.50 + 0.49/0.55 + 0.57/0.58 + 0.27/0.30. — Male. Carapace brownish, eyes surrounded with black. Sternum brownish. Labium, maxillae and chelicerae reddish-brown. Abdomen greyish-brown, with scutum. Spinnerets greyish-brown. Legs I brown with strong flat femora. Pedipalp structure shown in Fig. 4A–B. — Female. Carapace dark brown with black reticulate pattern, eyes surrounded by black rings. Whole surface of carapace with white adherent hairs. Abdomen grey with indistinct longitudinal streak. Legs I dark brown, remaining ones greyish-

brown. Epigyne with central pocket (Fig. 5A), its internal structures shown in Fig. 5B.

Habitat. Collected by sweeping in mixed woods, on meadows and dry places on the ground.

Distribution. Khabarovskii Krai and Primore.

Repeated study of specimens from Primore identified as *B. aurocinctus* by (Prószyński 1979 and Nenilin 1985) showed that they belonged to *H. nigriculus*.

Harmochirus pullus (Bösenberg et Strand, 1906)

Figs. 6–7

Material: Khabarovskii Krai, BR, 1♀, 18.VI.1987, D. Logunov (BI, No 667); Primore, Furugelm island, 1♂, 1♀ (incorrectly determined as "*Siler cupreus*"), 17–19.VII.1975, M. Shternbergs (BI, No 668, 669).

Description. Measurements (male/female). Cephalothorax: length 1.54/1.63, width 1.24/1.40, height 0.74/0.78. Abdomen: length 1.50/1.88, width 1.21/1.45. Eye field: length 0.93/1.03, AL




Fig. 6. *Harmochirus pullus* (Bösenberg et Strand), male, palpal organ. — A: ventral view. — B: lateral view.

1.03/1.15, PL 1.22/1.40. AM 0.36/0.36. Legs: I 1.03/0.90 + 0.59/0.50 + 0.81/0.58 + 0.53/0.53 + 0.39/0.35; II 0.69/0.73 + 0.38/0.45 + 0.41/0.40 + 0.37/0.43 + 0.30/0.30; III 0.88/0.93 + 0.41/0.48 + 0.44/0.50 + 0.44/0.53 + 0.32/0.35; IV 0.90/1.05 + 0.37/0.48 + 0.49/0.58 + 0.53/0.63 + 0.34/0.38. — Male. Colouration typical for the genus. Carapace convex uniform reddish-brown, surroundings of eyes black. Adherent white hairs on lateral surfaces of carapace. Sternum, labium, maxillae and chelicerae reddish-brown. Abdomen uniform dark grey with gleaming scutum. Spinnerets dark grey. Legs I: femora, patellae and tibiae reddish-brown, coxae, metatarsi and tarsi yellow. Femora with rows of scale-like black hairs on ventral surfaces, the same hairs on dorsal surfaces of tibiae (Fig. 7B). Remaining legs yellow. Pedipalp with long curved tibial apophysis (Fig. 6). — Female. Colouration resembling male. Abdomen without scutum. Young specimens yellow-greyish with white legs. Epigyne with large central depression and small pocket Fig. 7A).

Habitat. Collected on moist clearing with *Stratiotes* sp.

Distribution. New for Russia. Earlier known from Japan (Bösenberg & Strand 1906 and

Bohdanowicz & Prószyński 1987) and from Korea (Wesołowska 1981a and Paik 1987).

This species was recorded from Primore by Shternbergs (1988), but erroneously identified as *Siler cupreus* Simon.

Heliophanus dubius C. L. Koch, 1835

Material: Khabarovskii Krai, BR, 1♀, 14.VI.1987, D. Logunov (MNH).

Habitat. Collected by sweeping in deciduous forest.

Distribution. Widely distributed Palearctic species, new for Khabarovskii Krai. Recorded from the Russian Far East by Grube (1861), Dunin (1984), Nenilin (1985), Shternbergs (1988) and Marusik (1988).

Heliophanus lineiventris Simon, 1868

Material: Khabarovskii Krai, Nanaiskii distr., environs of Khabarovsk, 1♂, 15.VI.1931, V. Sychevskaya (ZMMU); the Amur river, Slavyanskii distr., 1♀, 20.VI.1989, D. Kurenschchikov (MNH), Amurskii distr., shore of Bolon lake, 51 km SW of Achat village, 3♀, 2.VI.1990, D. Kurenschchikov (IWP).




Fig. 7. *Harmochirus pullus* (Bösenberg et Strand), female. — A: epigyne. — B: first leg.

Habitat. Collected by sweeping in grass and shrubs.

Distribution. Widely distributed Palaearctic species, recorded from the Russian Far East by Kulczyński (1895), Prószyński (1976: map 74, 1979), Dunin (1985) and Nenilin (1985).

Heliophanus ussuricus Kulczyński, 1895

Material: Khabarovskii Krai, BR, 1♀, 5–17.VI.1987, D. Logunov (BI, No 498), 2♀ (MNH), 2♂, 2♀ (IBS); BR, the Chirka river, Akman village, 3♀, 15.VI.1989, S. Ivanov (BI, No 509).

Habitat. Collected by sweeping both in dry deciduous woods and in moister places (among sparse mosses and ferns in alder carr) as well as in clearings and on meadows.

Distribution. Eastern Palaearctic species, reported from the Russian Far East by Kulczyński (1895), Prószyński (1976: map 92, 1979), Dunin (1984), Nenilin (1985) and Marusik (1988).

Marpissa dersuuzalai sp. n.

Figs. 8–10

Material: Holotype: Khabarovskii Krai, BR, 1♂, 16–17.VI.1987, D. Logunov (BI, No 1119). — Paratypes: together with holotype, 6♂, 10♀ (BI, No 1120), 4♂, 3♀ (ZMMU, Ta-4665), 3♂, 4♀ (MNH, No 626); Amurskaya Oblast, Khinganskii reserve, 1♂, 15.VIII.1983, Yu. Marusik (ZMMU, Ta-4666). — Comparative material: Holotype of *Marpissa ibarakiensis*, Japan, Ibaraki, 1♀, 25.VI.1966, [leg.] Y. Suganami, coll. Yaginuma (IBS).

Diagnosis. The male in general appearance and colouration very similar to *Marpissa pulchra* Prószyński, *Marpissa magister* (Karsch), *Marpissa elongata* (Karsch) and *Marpissa nobilis* (Grube). May be easily told apart from them by bifurcated tibial apophysis (Fig. 8D). Similar, though not identical, apophysis possessed by “species 4” of Chikuni & Yaginuma (1976). The female closely resembles *Marpissa ibarakiensis* Bohdanowicz et Prószyński, but can be separated by configuration of spermathecae loops (cf. Figs. 10C and 11). For complete list of differences between females of *M. dersuuzalai* and *M. ibarakiensis* see Table 1. It is

Table 1. Differences between *Marpissa dersuuzalai* and *M. ibarakiensis*.

	<i>M. dersuuzalai</i>	<i>M. ibarakiensis</i>
Size	ca 7 mm	>10 mm
Cephalothorax length	ca 3 mm	>4 mm
Sternum	pale with dark margins	dark
Abdomen ventrally	3 thin lines	3 wide bands
Legs I	uniform	with contrasting Me, Ta
Insemination ducts	short, strongly coiled	long, slightly coiled




Fig. 8. *Marpissa dersuuzalai* sp. n., paratype, male. — A: general appearance. — B–D: palpal organ, ventral, dorsal and two lateral views.

very likely that *M. ibarakiensis* and the female of Chikuni & Yaginuma (1976), “species 4”, are conspecific.

Description. Measurements (male/female). Cephalothorax: length 2.53–2.88/2.85–3.02, width 1.80–2.05/1.95–2.30, height 0.85–1.03/0.93–1.20. Abdomen: length 2.88–3.45/2.85–3.02, width 1.45–1.70/1.95–2.30. Eye field: length

1.08–1.30/1.20–1.43, AL 1.33–1.48/1.40–1.63, PL 1.38–1.65/1.20–1.43. AM 0.40–0.43/0.47–0.50. Legs: I 1.63–1.95/1.55–2.00 + 0.88–1.20/0.95–1.15 + 1.38–1.68/1.20–1.45 + 0.98–1.2 5/0.85–1.05 + 0.53–0.63/0.45–0.55; II 1.15–1.33/1.20–1.45 + 0.65–0.73/0.75–0.90 + 0.73–0.83/0.70–0.95 + 0.70–0.85/0.65–0.90 + 0.43–0.48/0.40–0.50; III 1.20–1.40/1.25–1.50 + 0.65–0.73/




Fig. 9. *Marpissa dersuuzalai* sp. n., paratypes, female, epigyne.




Fig. 10. *Marpissa dersuuzalai* sp. n., paratypes, female. — A–C: epigyne and its internal structures. — D: abdominal pattern.




Fig. 11. *Marpissa ibarakiensis* Bohdanowicz et Prószyński, holotype (from Japan), female, internal structures of epigyne.

0.70–0.80 + 0.63–0.83/0.75–0.80 + 0.75–0.9 / 0.75–1.00 + 0.48–0.53/0.45–0.50; IV 1.28–1.63/1.50–1.80 + 0.68–0.83/0.75–1.00 + 0.98–1.20/1.05–1.30 + 0.88–1.0 / 1.00–1.20 + 0.50–0.53/0.50–0.55. — Male. Whole body elongated (Fig. 8A). Carapace dark brown with almost black punctured-reticulate eye field, behind eye field five spots formed by white hairs. Lateral edges of carapace with white narrow band. Sparse thin dark hairs on carapace. In vicinity of eyes few brown setae. Sternum dark brown. Clypeus brown, very low. Labium and maxillae dark brown with light margins. Chelicerae dark brown, promargin with two teeth, retromargin with single tooth. Abdomen black with bright golden shine and four pairs of white spots surrounded by black mat rings. Abdomen covered with thin dark hairs, denser at anterior margin. Ventrally abdomen dark with two longitudinal light lines, laterally additional lines formed by rows of points. Spinnerets dark. Legs brown with darker femora. Lateral surfaces of remaining segments of legs dark. Legs I thick. All legs with numerous thin dark hairs and brown spines. Pedipalp small, dark brown. Tibial apophysis two-cleft, pincers-shaped, posterior edge of cymbium with additional process, bulbous elongated, embolus geminate (Fig. 8B–D). — Female. Carapace brown with yellow or orange edges, eye field black; in

some specimens whole carapace black. Whitish-grey dense hairs cover carapace, near eyes sparse brown setae. Sternum yellow with dark margins. Labium and maxillae brown, sometimes with yellow margins. Chelicerae yellow or brown. Abdomen brown with golden shine, medially light streak formed by white hairs (Fig. 10D), in some specimens streak not visible. Ventrally abdomen light with three longitudinal dark stripes. Spinnerets brown. Legs yellow, in some specimens legs I brown. Pedipalp yellow. Epigyne small with two round copulatory openings (Figs. 9 and 10A–B). Insemination ducts rather short, weakly sclerotized, spermathecae multi-chambered (Fig. 10C).

Habitat. Collected by sweeping on meadows and wet places in clear woods (aspens, larch).

Distribution. Khabarovskii Krai and Amurskaya Oblast.

Etymology. Species named after Dersu Uzala, the legendary hero of V. A. Arsenev's novel, who supposedly died near the type locality.

Marpissa dybowskii (Kulczyński, 1895)

Material: Khabarovskii Krai, Evreiskaya Autonomna Oblast, Digun (?), 1♀, 13–18.VIII.1978, Belov (ZI); Primore, Ryazanovka, 1♀, VIII.1980, G. Belova (ZI).

Diagnosis. The species closely resembles *Marpissa pomatia* (Walckenaer). The main diagnostic differences are given in Prószyński (1979) and Bohdanowicz & Prószyński (1987).

Distribution. Eastern Palaearctic species, new for Khabarovskii Krai. Previously recorded from Primore by Prószyński (1976: map 118, 1979), Dunin (1984), Nenilin (1985) and Shternbergs (1988).

Marpissa elongata (Karsch, 1879)

Material: Khabarovskii Krai, environs of Khabarovsk, the Amur river, Zelonyi island, 1♂, 7.VII.1931, V. Sychevskaya (ZMMU); Amurskaya Oblast, Khinganskii reserve, Lebedinskoe forest-range, 1♂, 2♀, 8.VIII.1983, Yu. Marusik (ZI); Lebedinskoe lake, 4♂, 2–3.VIII.1991, D. Kurenshchikov (BI, No 1121); Primore, Chernigovskii distr., Dmitrevka, 1♂, 16.VII.1989, A. Borok (BI, No 1122); Prikhankaiskii distr., Khazan lake, Kamen-Rybolov village, 2♂, 2♀, 4.V–24.VII.1908, A. Cherskii (ZI).




Fig. 12. *Marpissa magister* (Karsch), male. — A–C: palp organ, ventral, lateral and dorsal views. — D: lateral view of tibial apophysis.

Diagnosis. The species resembles *Marpissa nobilis* (Grube) and *Marpissa pulchra* Prószyński. The diagnostic features are given in Prószyński (1973), Chikuni & Yaginuma (1976), Wesołowska (1981a), as well as Bohdanowicz & Prószyński (1987).

Habitat. Collected by sweeping grass.

Distribution. Eastern Palaearctic species, new for Khabarovskii Krai. From the Russian Far East reported from Primore (Prószyński 1976: map 121, 1979, Dunin 1984, Nenilin 1985, and Shternbergs 1987).

Marpissa magister (Karsch, 1879)

Figs. 12–13

Material: Khabarovskii Krai, BR, 1♀, 19.VI.1987, D. Logunov (BI, No 1123), 1♂, 17.VII. 1987, D. Logunov (MNH); Nizhnespaskoe, 53 km SE of Khabarovsk, 1♂, 15.VI.1988, D. Kurenshchikov (BI, No 1124); Primore, Khazanskii distr., Talmi lake, Khazan village, 1♀, 27.VII. 1976, B. Zakharov (BI, No 1125).

Diagnosis. From other similar *Marpissa* species the male is told apart by broader tibial apophysis (Fig. 12 B and 12 D), the female by shape of epigyne and spermathecae (Fig. 13 C, see also Chikuni & Yaginuma 1976 and Bohdanowicz & Prószyński 1987).

Description. Measurements (male/female). Cephalothorax: length 2.93/2.55, width 2.00/1.73, height 0.98/0.78. Abdomen: length 3.65/3.40, width 1.75/1.45. Eye field: no measurements. AM 0.46/0.49. Legs: I 2.25/1.45 + 1.15/0.85 + 1.60/1.15 + 1.10/0.80 + 0.55/0.45; II 1.25/1.00 + 0.75/0.65 + 0.75/0.65 + 0.65/0.60 + 0.45/0.40; III 1.25/1.00 + 0.60/0.65 + 0.70/0.55 + 0.70/0.60 + 0.45/0.45; IV 1.60/1.35 + 0.75/0.75 + 1.10/0.95 + 0.95/0.85 + 0.45/0.45. — Male. Whole body coloured dark brown. White hairs form separate spots on carapace and abdomen (Fig. 13A). At anterior margin of abdomen narrow white band. Sternum black, labium and maxillae black with yellow margins. Legs brown with yellow trochanters and tarsi. Pedipalp shown in Fig. 12. Tibial apophysis broad (Fig. 12B and 12D).




Fig. 13. *Marpissa magister* (Karsch), male and female. — A: general appearance of male. — B: general appearance of female. — C: epigyne.

— Female. Carapace light brown with darker eye field, eyes surrounded with black. Sternum centrally yellow, marginally brown. Labium grey with lighter margin, maxillae yellow, chelicerae orange. Abdomen with two longitudinal broad brown stripes with golden shine and four pairs of square black spots (Fig. 13B). Ventrally abdomen yellowish brown with three longitudinal brown streaks. Spinnerets yellow with dark dots. Legs yellowish-brown, legs I a little darker. Epigyne small with depression in posterior part, divided by triangular median ridge (Fig. 13C).

Habitat. Collected by sweeping in dry deciduous woods and meadows, as well as in moister places.

Distribution. In the Russian Far East: Khabarovskii Krai and Primore (Dunin 1984 — incorrectly determined as “*M. pichoni*”, Marusik & Logunov in press). Earlier known from Japan, Viet-Nam and China (after Žabka 1985).

Dunin’s (1984) records of *Marpissa pichoni* (Schenkel) from Primore actually refer to *M. magister*, as can clearly be seen from his figures (42–44).

Marpissa nobilis (Grube, 1861)

Figs. 14–15

Material: Khabarovskii Krai, BR, 1♂, 1♀, 16–17.VI.1985, D. Logunov (BI, No 1126, 1127); Amurskaya Oblast, Khinganskii reserve, 3♂, 1–24.VIII.1983, Yu. Marusik (ZMMU), 3♂ (ZI); the Ussuri river, 1♂ lectotype, 1♂ paralectotype, [leg.] Schrenk, coll. Grube (MNH, No 436).

Diagnosis. The species very close to *Marpissa pulchra* Prószyński. The male can be separated by lack of tooth on tibial apophysis (cf. Figs. 14B–E and 16A–C) and by colouration of legs. Patellae and tibiae II–IV of *M. nobilis* are yellow, whereas those of *M. pulchra* are brown. The female may be easily recognized by structure of vulva (cf. Figs. 15C and 17C).

Description. Measurements (male/female). Cephalothorax: length 3.13/3.03, width 2.20/2.03, height 1.05/0.98. Abdomen: length 3.90/5.25, width 1.80/2.25. Eye field: length 1.38/1.28, AL 1.55/1.48, PL 1.55/1.55. AM 0.54/0.41. Legs: I 3.20/1.70 + 1.60/1.10 + 2.90/1.50 + 1.90/1.05 + 0.50/0.50; II 1.55/1.25 + 0.85/0.95 + 1.10/0.80 + 0.90/0.75 + 0.50/0.40; III 1.50/1.35 + 0.80/0.75




Fig. 14. *Marpissa nobilis* (Grube), male. — A: general appearance. — B–D: tibial apophysis, ventral, dorsal and lateral views. — E: paralectotype, lateral view of tibial apophysis.

+ 0.95/0.75 + 1.00/0.85 + 0.55/0.55; IV 1.90/1.60 + 0.85/1.25 + 1.50/0.85 + 1.30/1.15 + 0.55/0.45. — Male. Whole body coloured dark brown, eyes surrounded by black. Carapace and abdomen with ornament composed of white hair spots (Fig. 14A). Abdomen ventrally with two thin light longitudinal lines. Legs I dark brown with yellowish tarsi. Remaining legs yellow, only femora brown with yellow streaks. Pedipalp structure similar to *M. pulchra*, details of structure of tibial apophysis shown in Fig. 14B–E. — Female. Carapace brown with orange margins, eye field black. Sternum yellowish-brown. Labium and chelicerae brown, maxillae yellow. Abdomen yellowish with two broad dark longitudinal bands (Fig. 15A), ventrally with three thin brown lines. Spinnerets yellow. All legs yellow, only patella, tibia and metatarsus of leg I brownish. Epigyne with small depression in pos-

terior part, partially divided by wide median ridge (Fig. 15B). Its internal structures as in Fig. 15C.

Habitat. Collected by sweeping on meadows and in swampy woods.

Distribution. Eastern Palaearctic species, reported from the Russian Far East (Grube 1861, Prószyński 1971, 1979) and China (Schenkel 1963 and Wesołowska 1981b).

Marpissa pomatia (Walckenaer, 1802)

Material: Khabarovskii Krai, BR, 3♂, 12♀, 14–15.VI.1987, D. Logunov (BI, No 896, 1128), 2♂, 4♀ (MNH); BR, Bychikha village, 2♀, 15.V.1988, S. Ivanov (BI, No 891); BR, the Chirka river, 3♀, 26–31.V.1988, S. Ivanov (BI, No 882, 892); Ulchskii distr., the Asyat river, 1♀, 25.VI.1990, D. Kurenshchikov (IWP); Amurskaya Oblast, Selezdzhinsk, 1♂, VI–VII.1985, A. Kudryavtsev (ZMMU).




Fig. 15. *Marpissa nobilis* (Grube), female. — A: general appearance. — B–C: epigyne and its internal structures.

Diagnosis. Resembles *Marpissa dybowskii* Kulczyński. Good diagnostic figures in Prószyński (1979: figs. 188–199 only).

Habitat. Collected by sweeping in deciduous woods and in clearings.

Distribution. Widely distributed Palearctic species. Recorded from the Russian Far East by Kulczyński (1926), Prószyński (1976: map 114, 1979), Azheganova & Stenchenko (1977), Dunin (1984) and Nenilin (1985).

***Marpissa pulchra* Prószyński, 1976**

Figs. 16–17

Material: Khabarovskii Krai, BR, 1♂, 6♀, 16–19.VI.1987, D. Logunov (BI, No 1129, 1131), 2♀ (MNH); Amurskaya Oblast, Arkhara village, 1♂, 24.VIII.1983, Yu. Marusik (BI, No 1132).

Diagnosis. See under *Marpissa nobilis*. Good diagnosis of *M. pulchra* is given in Chikuni (1989: figs. 14–17).

Description. Measurements (male/female). Cephalothorax: length 2.53/2.88–3.00, width

1.70/1.95–2.05, height 0.88/0.80–1.00. Abdomen: length 3.45/5.30, width 1.40/2.15. Eye field: length 1.10/1.28–1.38, AL 1.28/1.30–1.80, PL 1.28/1.45–1.80. AM 0.42/0.46–0.50. Legs: I 1.85/1.80–2.00 + 1.15/1.00–1.20 + 1.80/1.60–1.80 + 1.05/1.05–1.25 + 0.60/0.60; II 1.20/1.15–1.35 + 0.65/0.70–0.75 + 0.75/0.75–0.85 + 0.65/0.65–0.80 + 0.40/0.50; III 1.05/1.20–1.35 + 0.60/0.60–0.65 + 0.70/0.65–0.75 + 0.70/0.80–0.85 + 0.50/0.45–0.55; IV 1.20/1.50–1.65 + 0.70/0.80–0.85 + 1.05/1.15–1.20 + 0.95/1.00–1.15 + 0.50/0.50–0.55. — Male. Carapace dark brown, eye field black, behind eye field two white spots. Sternum and chelicerae dark brown, labium and maxillae brown with yellow margins. Abdomen brown with golden gleam, with scutum. Four pairs of white spots, composed of hairs, on abdomen (Fig. 16D). Spinnerets brown. All legs dark brown, only tarsi yellowish. Pedipalp structure shown in Fig. 16A–C, characteristic small tooth on apex of tibial apophysis. — Female. Carapace brown, covered with white hairs, eye field black. Abdomen grey with medial longitudinal broad white band and four pairs of black spots (Fig. 17D).




Fig. 16. *Marpissa pulchra* Prószyński, male. — A–C: palpal organ, ventral, lateral and dorsal views. — D: general appearance.

Sides of abdomen yellow, ventrally abdomen with wide longitudinal brown band and two thin yellow lines. Legs I brown with yellow metatarsi and tarsi, remaining ones yellow with brownish proximal parts of tarsi. Pedipalp yellow with brownish tarsus. Epigyne and its internal structures shown in Fig. 17A–C.

Habitat. Collected on grass in paludal forest.

Distribution. New for Russia. Earlier known from Japan (Prószyński 1976, Chikuni & Yaginuma 1976 — as “species 3”, Chikuni 1989, Bohdanowicz & Prószyński 1987) and Korea (Wesołowska 1981a).

Marpissa pulla (Karsch, 1879)

Material: Khabarovskii Krai, BR, 1♀, 16.VI.1987, D. Logunov (BI, No 1133), 1♀, 19.VI.1987 (MNH); Primore, Sadgorod, 1♀, 13.VI.1978, Belov (ZI).

Diagnosis. The species similar to *Marpissa dybowskii* Kulczyński and *Marpissa pomatia* (Walckenaer). Good diagnostic figures in Weso-

łowska (1981b) and Bohdanowicz & Prószyński (1987).

Habitat. Collected by sweeping grass on forest clearing.

Distribution. Eastern Palearctic species, new for Khabarovskii Krai. Previously recorded in the Russian Far East from Primore by Prószyński (1979 — incorrectly determined as “*M. pomatia*”), Nenilin (1985) and from Furugelm island by Shternbergs (1988).

Among specimens treated by Prószyński (1979) as *M. pomatia*, there were also individuals of *M. pulla*, as is clearly visible in his figures (184–187).

Marpissa zebra sp. n.

Fig. 18

Material: Holotype: Khabarovskii Krai, BR, 1♂, 17.VI.1987, D. Logunov (BI, No 1134). — Paratype: Amurskaya Oblast, Khinganskii reserve, 1♂, 4.VIII.1983, Yu. Marusik (ZI).




Fig. 17. *Marpissa pulchra* Prószyński, female. — A–C: epigyne and its internal structures. — D: general appearance.

Diagnosis. The species can be easily distinguished from other *Marpissa* species by its very characteristic colouration pattern (Fig. 18C) and small size. Palpal structure very similar to *Marpissa elongata* (Karsch), *Marpissa nobilis* (Grube) and *Marpissa pulchra* Prószyński, but may be separated by yellow colouration of cymbium and configuration of tibial apophysis and embolus.

Description. Measurements (holotype). Cephalothorax: length 1.98, width 1.35, height 0.65. Abdomen: length 2.53, width 1.02. Eye field: length 0.90, AL 1.00, PL 1.05. AM 0.60. Legs: I 1.18 + 0.70 + 1.05 + 0.85 + 0.43; II 0.83 + 0.35 + 0.55 + 0.50 + 0.30; III 0.80 + 0.40 + 0.50 + 0.50 + 0.33; IV 0.95 + 0.53 + 0.73 + 0.65 + 0.35. — Male. Carapace yellow with brown reticulate pattern, near eyes black. Sternum, maxillae and chelicerae yellow, labium brown with light margin. Abdomen yellow with four transversal brown streaks and narrow median line along anterior third of its length (Fig. 18C).

Ventrally abdomen light yellow with three longitudinal dark streaks. Legs I yellow with darker metatarsi and tarsi, remaining legs yellow with longitudinal dark lines, on tibiae elongated brown spots. Pedipalp shown in Fig. 18A–B. — Female unknown.

Habitat. Collected from wet places in mixed woods and on meadows.

Distribution. Khabarovskii Krai and Amurskaya Oblast.

Myrmarachne formicaria (De Geer, 1778)

Figs. 19–21

Material: Khabarovskii Krai, BR, 1♂, 1♀, 16.VI.1987, D. Logunov (BI, No 901, 902); Nanaiskii distr., 1♂, 1983, N. Ryabinin (ZMMU); Amurskaya Oblast, Khinganskii reserve, Antonovskoe forest-range, 1♀, 2–3.VIII.1983, Yu. Marusik (ZI); the Ussuri river, 1♀, 10.VII.1931, V. Sychevskaya (ZMMU, Ta-3427); Primore, Ussuriiskii reserve, 1♀, 5.VIII.1975, M. Shternbergs (BI, No 903).




Fig. 18. *Marpissa zebra* sp. n., holotype, male. — A–B: palpal organ, ventral and lateral views. — C: general appearance.

Diagnosis. See under *Myrmarachne lugubris* and Table 2.

Description. Measurements (male/female). Cephalothorax: length 1.63/2.50, width 0.88/1.08, height 0.63/0.80. Abdomen: no measurements. Eye field: length 0.80/1.00, AL 0.83/1.05, PL 0.95/1.18. AM 0.23/0.36. Legs: I 0.98/1.18 + 0.45/0.63 + 0.80/0.95 + 0.50/0.55 + 0.33/0.35; II 0.75/0.80 + 0.38/0.50 + 0.48/0.68 + 0.38/0.50 + 0.30/0.30; III 0.75/0.80 + 0.35/0.45 + 0.68/0.50 + 0.30/0.53 + 0.30/0.30; IV 1.00/1.40 + 0.43/0.63 + 0.93/1.20 + 0.78/1.13 + 0.35/0.40. — Male. Dorsal aspect in Fig. 19A. Carapace reddish with darker punctured-reticulate eye field, eyes surrounded by black. Labium and maxillae dark brown, sternum lighter. Chelicerae very elongated, horizontal, dark brown, with numerous teeth (Fig. 19B). Cheliceral fang with small protuberance on ventro-lateral side (Fig. 19C). Abdomen elongated brown, posteriorly darker, with anterior and posterior scutum, whole covered by sparse adherent hairs. Few white hairs form two small transverse spots mid-length of posterior scutum, posterior part of this scutum almost black.




Fig. 19. *Myrmarachne formicaria* (De Geer), male. — A: general appearance. — B: cheliceral dentition. — C: fang of chelicerae, ventro-lateral view.




Fig. 20. *Myrmarachne formicaria* (De Geer), male and female. — A–C: palpal organ, ventral, lateral and dorsal views. D–E: epigyne and its internal structures.

Ventrally abdomen light with longitudinal broad grey streak medially and many dark lines laterally. Spinnerets brown. Legs I yellow with brown femora and metatarsi. Remaining legs yellowish. Pedipalp brown, small with distinctive tibial apophysis (Fig. 20A–C). — Female. Colouration as in male, but darker. Carapace reddish with

black eye field. Sternum dark brown. Labium and maxillae brown with yellow margins, chelicerae brown. Abdomen grey with two pairs of white small transversal spots dorsally, on lateral surfaces of abdomen white stripes. Coxae yellow, all of them — except first — with broad dark stripes. Legs I shown in Fig. 21A. Legs III and

Table 2. Differences between *Myrmarachne lugubris* and *M. formicaria*.

		<i>M. lugubris</i>	<i>M. formicaria</i>
Male and female	Carapace	uniform — blackish	reddish, eye field black
	Tarsus I	brown (Fig. 21B)	yellow (Fig. 21A)
	Coxae III–IV	brownish	yellow
Male only	Cheliceral fang	without teeth	with additional tooth (Fig. 19C)
	Cymbium	with tuft of hairs (Fig. 22A)	hairs absent (Fig. 20A)
	Scutae	both brownish	first and frontal part of second reddish
Female only	Abdomen	uniform — dark grey	with pair of transversal light spots
	Epigyne	Fig. 23A–B	Fig. 20D




Fig. 21. *Myrmarachne formicaria* (De Geer) and *Myrmarachne lugubris* (Kulczyński). — A: first leg of female of *M. formicaria*. — B: first leg of female of *M. lugubris*. — C–D: *M. lugubris*, cheliceral dentition of male and female.

IV with black longitudinal streaks on patellae and tibiae. Epigyne with large, more or less triangular depression, partially divided by median ridge (Fig. 20D). Internal structures of epigyne shown in Fig. 20E.

Habitat. Collected in wet woods (oak, birch, ferns) in litter.

Distribution. Palearctic species displaying Euro-Far Eastern disjunction of the range. New for Khabarovskii Krai. Recorded from the Russian Far East by Prószyński (1976: map 197, 1979), Azheganova & Stenchenko (1977), Dunin (1984), Oliger (1984) and Nenilin (1985).

Myrmarachne lugubris Kulczyński, 1895

Figs. 21–23

Material: Khabarovskii Krai, BR, 1♂, 1♀, 14.VI.1987, D. Logunov (MNH); BR, the Chirka river, 1♂, 1♀, 17–30.V.1988, S. Ivanov (BI, No 1136, 1137); Khabarovsk,

cemetery, 25.VI.1931, 2♀, V. Sychevskaya (ZMMU); Nainiskii distr., Slavyanka, 1♂, 23.V.1988, D. Kurenschikov (BI, No 1135), 2♀ (ZMMU); Primore, Vladivostok, 1♂, 1♀, 1.VI.1979, B. Zakharov (ZI).

Diagnosis. Very similar to *Myrmarachne formicaria* (De Geer), therefore often mistaken with this species. Complete list of differences between those species is presented in Table 2. *M. formicaria* and *M. lugubris* occur in different microhabitats. In Bolshekhekhtsirsky reserve the former species was found in forest litter and the latter was recorded while sweeping from grasses in woodland glades.

Description. Measurements (male/female). Cephalothorax: length 2.75/2.95, width 1.33/1.43, height 0.95/1.08. Abdomen: no measurements. Eye field: length 1.13/1.18, AL 1.18/1.28, PL 1.28/1.35. AM 0.38/0.35. Legs: I 1.75/1.55 + 0.73/0.68 + 1.53/1.35 + 1.03/0.85 + 0.55/0.45; II 1.28/1.23 + 0.65/0.55 + 0.98/0.90 + 0.75/0.75 + 0.40/0.40; III 1.43/1.40 + 0.63/0.60 + 1.05/




Fig. 22. *Myrmarachne lugubris* (Kulczyński), male, palpal organ. — A: ventral view. — B: lateral view. — C: dorsal view.




Fig. 23. *Myrmarachne lugubris* (Kulczyński), female. — A–B: epigyne. — C: internal structures of epigyne.

1.03 + 1.08/1.10 + 0.43/0.40; IV 2.15/2.15 + 0.68/0.80 + 1.60/1.73 + 1.50/1.63 + 0.50/0.48. — Male. Ant-like spider. Carapace brownish-

black. Carapace with constriction behind eye field. Clypeus very low. Sternum dark brown, labium and maxillae brown with yellow margins.




Fig. 24. *Phintella arenicolor* (Grube), lectotype, male. — A–B: palpal organ, ventral and lateral views. — C: tibial apophysis.

Chelicerae yellowish-brown, very long and slender, horizontal (Fig. 21C). Abdomen dark grey. Spinnerets dark. Pedipalp structure shown in Fig. 22. Characteristic tuft of hairs on basal part of cymbium (Fig. 22A). — Female. Colouration resembling male. Coxae I–II light yellow, III–IV dark brown. Legs I shown in Fig. 21B, legs II yellow with dark stripes on femora, tibiae and patellae, legs III brown with yellow metatarsi and tarsi, legs IV brown. Epigyne with great incision on posterior edge. Large depression partially divided by median ridge (Fig. 23A–B). Internal structures of epigyne as in Fig. 23C.

Habitat. Collected by sweeping in deciduous forest (aspen, birch, oak). Also recorded in stream submersion areas.

Distribution. Eastern Palearctic species, new for Khabarovskii Krai. From the Russian Far East reported by Prószyński (1976: map 202, 1979), Dunin (1984), Oligier (1984) and Nenilin (1985).

Phintella arenicolor (Grube, 1861)

Figs. 24–27

Phintella mellotei (Simon, 1888), syn. n.

Material: Khabarovskii Krai, BR, 2♂, 17.VI.1987, D. Logunov (BI, No 684); Nanaiskii distr., Slavyanka, 1♀, 26.V.1989, D. Kurenschikov (BI, No 683); environs of Khabarovsk, the Amur river, Zelonyi island, 1♀, 22.VI.1931, V. Sychevskaya (ZMMU); the Amur river, 1♂ lectotype of *Attus arenicolor* (nomen museorum “*Salticus pallidus*”), 2 juv. ♀, [leg.] Dittmar, coll. Grube (MNH, No 429); Amurskaya Oblast, Khinganskii reserve, Antonovskoe forest-range, 2♀, 1.VIII.1983, Yu. Marusik (ZI); Primore, Khasanskii distr., Kedrovaya Pad reserve, 1♀, 29.VIII.1968, F. Popov (ZI); Primorskaya station, 1♂, VII.1979, B. Zakharov (BI, No 682); Chuguevskii distr., 1♀, 23.IX.1974, (ZMMU); Ussuriiskii reserve, 1♀, 3.X.1977, G. Kurcheva, E. Mikhaleva (ZMMU); middle stream of the Bikin river, 1♀, VIII.1978, Shibnev (BI, No 685); Prikhankaikskii distr., Khorol village, 2♀, 18.VII.1968, F. Popov (ZI); Khazan lake, Kamen-Rybolov village, 1♂, 1♀, 19.VII.1968, F. Popov (ZI); Furugelm island, 1♂, 18.VIII.1975, M. Shternbergs (BI, No 686). — Additional material: Japan, Goto Islands, Nagasaki Pref., 2♂ (incorrectly determined as “*P. castrisiana*”), 14–20.V.1968, coll. Yaginuma (IBS).




Fig. 25. *Phintella arenicolor* (Grube), male, palpal organ. — A: ventral view. — B: lateral view.

Diagnosis. In colouration similar to *Phintella parva* (Wesołowska), but differs in absence of dark longitudinal lines on femora III and IV, smaller size and shape of tibial apophysis. In structure of genitalia *P. arenicolor* is very similar to *Phintella castriesiana* (Grube). The male can be distinguished by structure of tibial apophysis (cf. Fig. 27C and 27D), thinner apical part of tegulum (cf. Fig. 27E and 27F) and body colouration (cf. Fig. 27A and 27B). Distinguishing the females of both species is especially difficult. The female may be recognized by shape and smaller size of spermathecae (cf. Figs. 26C and 29B–C).

Description. Measurements (male/female). Cephalothorax: length 1.50/1.68, width 1.15/1.28, height 0.75/0.75. Abdomen: length 1.63/2.50, width 0.93/1.73. Eye field: length 0.90/0.88, AL 0.98/1.03, PL 0.93/1.03. AM 0.30/0.30. Legs: I 1.10/1.90 + 0.63/0.53 + 0.88/0.58 + 0.73/0.78 + 0.43/0.35; II 0.98/0.88 + 0.48/0.45 + 0.65/0.53 + 0.70/0.55 + 0.38/0.30; III 0.98/1.00 + 0.43/0.45 + 0.75/0.63 + 0.90/0.78 + 0.40/0.38; IV 1.15/1.35 + 0.43/0.48 + 0.88/0.85 + 0.95/0.95 + 0.43/0.40. — Male. Carapace light yellow, eyes surrounded by black. Carapace covered with translucent scales, on its posterior part scales brown. Sternum, labium and maxillae yellow, chelicerae yellow with dark longitudinal stripes on front.




Fig. 26. *Phintella arenicolor* (Grube), female. — A: general appearance. — B–C: epigyne and its internal structures.

Abdomen yellow with indistinct darker ornament (Fig. 27A). Spinnerets yellow. Legs yellow with dark longitudinal stripes on femora I–II and brown rings on apical ends of tibiae I–II. Pedipalp structure in Figs. 24 and 25. — Female. Dorsal aspect in Fig. 26A. Colouration similar to male, but chelicerae and legs uniform yellow. Epigyne weakly sclerotized (Fig. 26B), insemination ducts rather straight, spermathecae more or less spherical (Fig. 26C).

Habitat. Collected in swampy woods and on meadows in grass.

Distribution. Eastern Palearctic species, new for Khabarovskii Krai. Previously recorded from the Russian Far East as *Phintella difficilis* (Bösenberg et Strand) by Prószyński (1976: map 103, 1979: figs. 143–144 only), Dunin (1984) and Shternbergs (1988).

P. arenicolor was described by Grube (1861), and was later synonymized with *P. castriesiana* (Grube) by Prószyński (1979). However, a renewed examination of the type material leads us to the conclusion that *P. arenicolor* constitutes a valid species, and that high variability of colouration and of reproductive structures recorded earlier in *P. castriesiana* (Prószyński 1979 and Matsumoto 1989) could be due to the fact that they were unaware that they were actually dealing with two species.




Fig. 27. *Phintella arenicolor* (Grube) (left) and *Phintella castriesiana* (Grube) (right), males. — A–B: general appearance. — C–D: lateral view of tibial apophysis. — E–F: embolus.

P. arenicolor is new for Japan; by Bohdanowicz & Prószyński (1987) erroneously reported as *Phintella castriesiana* (Grube).

Phintella castriesiana (Grube, 1861)

Figs. 27–29

Material: Neotype: Khabarovskii Krai, BR, 1♂, 5–22.VI.1987, D. Logunov (MNH, No 625). Together with neotype, 33♂, 45♀ (BI, No 648, 650, 652, 654, 655, 656, 662, 663), 6♂, 9♀ (MNH); the Khor river, 17 km upper of Kutuzovka village, 1♂, 6–10.VI.1985, S. Pustoroi, A. Makoedov (ZMMU).

Designation of the neotype. *Phintella castriesiana* was described (Grube 1861) on the basis of a single female. The holotype (examined) kept in collection of Museum of Natural History in Wrocław consists now only of separated cephalothorax, devoided of legs and pedipalps (see also Prószyński 1971); it is thus devoid of diagnostic features. The designation of the neotype is necessary in this case, as there are serious problems with differentiating between *P. castriesiana* and the closely related *P. arenicolor* (a list of features differentiating these species is given under “diagnosis” in description of *P. arenicolor*). *P. castriesiana* is relatively well known. Hence, there is no doubt that the neotype

is consistent with the holotype. The type locality was not given precisely by Grube (1861) but it is known that the holotype was collected in the Amur river valley, between the mouth of the Ussuri river and Nikolaevsk na Amure. The Bolshekhkhechtsirskii reserve, from where the neotype originates is situated in the SW part of this area.

Diagnosis. The largest and the darkest species of the genus *Phintella* from the Russian Far East. In the genitalia structure very close related to *Phintella arenicolor* (Grube) (differentiation see under *P. arenicolor*).

Description. Measurements (male/female). Cephalothorax: length 1.95–2.30/2.08–2.33, width 1.45–1.65/1.53–1.70, height 0.88–1.03/0.85–0.95. Abdomen: length 2.00–2.35/2.95–3.38, width 1.15–1.38/2.00–2.08. Eye field: length 1.00–1.08/1.03–1.23, AL 1.23–1.28/1.23–1.44, PL 1.18–1.25/1.20–1.38. AM 0.40/0.38–0.45. Legs: I 1.38–1.60/1.10–1.33 + 0.68–0.78/0.60–0.75 + 1.03–1.23/0.80–0.88 + 0.95–1.0 3/0.68–0.80 + 0.50–0.55/0.43–0.48; II 1.25–1.38/1.08–1.23 + 0.60–0.65/0.53–0.60 + 0.88–0.93/0.70–0.80 + 0.90–0.9 5/0.68–0.73 + 0.43–0.45/0.38–0.45; III 1.33–1.40/1.23–1.33 + 0.58–0.63/0.55–0.58 + 0.88–1.03/0.75–0.93 + 1.15–1.25/0.98–1.10 + 0.45/0.48–0.50; IV 1.40–1.68/1.38–1.58 + 0.60–0.65/0.55–0.68 + 1.10–1.25/1.00–1.18 + 1.35–1.3 8/1.18–1.35 + 0.48–0.50/0.45–0.53. — Male.




Fig. 28. *Phintella castriesiana* (Grube), male, palpal organ. — A: ventral view. — B: lateral view.

Carapace yellowish-brown with black veins, eyes surrounded by black. In vicinity of eyes white scales. Sternum, labium and maxillae yellow, chelicerae brownish. Abdomen yellow with dark grey pattern (Fig. 27B), ventrally light with dark longitudinal streak. Spinnerets yellowish-grey. Legs yellow with brown longitudinal stripes on femora I–III and brown dots on proximal and distal parts of tibiae I–II. Pedipalp structure shown in Fig. 28. — Female. Colouration resembling male but slightly paler. All legs yellow. Epigyne and its internal structures shown in Fig. 29.

Habitat. Collected on grass in mixed woods.

Distribution. Eastern Palaearctic species, recorded from the Russian Far East by Grube (1861), Kulczyński (1895), Prószyński (1971, 1976: map 95, 1979), Dunin (1984) and Nenilin (1985). Records from Western Palaearctic refer most probably to another, closely related species (this problem will be dealt with separately).

Phintella linea (Karsch, 1879)

Figs. 30–31

Material: Khabarovskii Krai, BR, 1♂, 1♀, 17.VI.1987, D. Logunov (BI, No 558, 559), 1♀ (MNH).

Diagnosis. The male can be easily distinguished by shape of tibial apophysis, great sharp embolus (Fig. 30A–B) and contrasting colouration (Fig. 30C). The female may be separated by the spermathecae being larger than in other *Phintella* species (Fig. 31C).

Description. Measurements (male/female). Cephalothorax: length 1.53/1.70–1.80, width 1.18/1.30–1.38, height 0.78/0.83–0.85. Abdomen: length 1.25/2.05–3.25, width 1.88/1.43–2.38. Eye field: length 0.88/0.95–1.00, AL 1.05/1.15–1.18, PL 1.03/1.20. AM 0.33/0.35–0.38. Legs: I 0.93/0.95–1.00 + 0.45/0.50–0.53 + 0.65/0.65–0.68 + 0.55/0.53–0.58 + 0.35/0.35–0.38; II 0.83/0.95 + 0.38/0.45–0.48 + 0.55/0.55–0.58 + 0.53/0.53–0.55 + 0.33/0.30–0.33; III 0.95/1.00–1.03 + 0.42/0.43–0.53 + 0.55/0.60–0.63 + 0.68/0.75–0.78 + 0.40/0.38; IV 1.03/1.20–1.28 + 0.43/0.48–0.53 + 0.65/0.75–0.83 + 0.80/0.95–1.00 + 0.40/0.35–0.38. — Male. Dorsal aspect as in Fig. 30C. Carapace brownish-yellow with dark brown spots. On carapace translucent scales with green shine. Sternum yellow with brown margin. Clypeus, labium and chelicerae brown. Maxillae yellow. Abdomen yellow with brown pattern composed of longitudinal and diagonal stripes, ventrally yellow with single longitudinal brown band. Spinnerets greyish. Legs yellow with brown rings




Fig. 29. *Phintella castrisiana* (Grube), female. — A: epigyne. — B–C: internal structures of epigyne.

on segment ends. Pedipalp yellow with brown femur and basal part of cymbium. Palpal structure shown in Fig. 30A–B. — Female. Dorsal aspect in Fig. 31A. Carapace yellow, eyes surrounded by black. Sternum yellow with brown margin. Labium, maxillae, chelicerae, pedipalps and legs yellow. Abdomen as in male. Epigyne as in Fig. 31B. Insemination ducts short, spermathecae large, oval (Fig. 31C).

Habitat. Collected by sweeping grass in wet woods.

Distribution. Eastern Palaearctic species, new for Russia. The earlier record (Nenilin 1985) was based on misidentification. Reanalysis of his material showed that these specimens actually belonged to *Phintella popovi* Prószyński. The record from Korea (Wesołowska 1981a) is also

erroneous. The pictures in her paper actually refer to the female of *P. popovi* (figs. 37–38) and the male of *P. arenicolor* (figs. 42–44). Thus, so far, *P. linea* has been doubtless recorded only from Japan (Karsch 1879, Bösenberg & Strand 1906, Prószyński 1973, 1976: map 102 — part, 1978, Bohdanowicz & Prószyński 1987 and Matsumoto 1989).

Phintella parva (Wesołowska, 1981)

Figs. 32–33

Material: Khabarovskii Krai, BR, 6♂, 2♀, 2–14.VI.1987, D. Logunov (BI, No 601, 603, 604, 605), 2♂, 4♀ (MNH); BR, the Chirka river, 1♀, 15.VI.1989, S. Ivanov (BI, No 602); Primore, Chuguevskii distr., 1♀, 9.IX.1974, G. Kurcheva




Fig. 30. *Phintella linea* (Karsch), male. — A–B: palpal organ, ventral and lateral views. — C: general appearance.




Fig. 31. *Phintella linea* (Karsch), female. — A: general appearance. — B–C: epigyne and its internal structures.

(ZMMU); Ussuriiskii reserve, 1♀, 5.VIII.1975, M. Shternbergs (BI, No 600). — Additional material: Japan, Mie Pref., 2♀ (incorrectly determined as "*P. mellotei*"), 28.V.1967, coll. Yaginuma (IBS).

Diagnosis. Resembles *Phintella popovi* (Prószyński); the male may be easily separated by straight tibial apophysis and curved embolus (cf. Figs. 32 and 34A–B), the female by longer insemination ducts (cf. Figs. 33C and 34D).

Description. Measurements (male/female). Cephalothorax: length 1.78–2.05/1.73–1.90, width 1.28–1.55/1.25–1.38, height 0.95/0.73–0.88. Abdomen: length 1.83–2.00/2.88–3.08, width 1.05–1.20/1.80–2.18. Eye field: length 0.83–1.05/0.88–1.05, AL 1.03–1.20/1.08–1.20, PL 0.98–1.15/1.08–1.20. AM 0.30–0.38/0.35–0.38. Legs: I 1.20–1.43/0.90–1.03 + 0.70–0.83/0.45–0.53 + 0.95–1.18/0.60–0.73 + 0.80–0.9 5/0.53–0.60 + 0.43–0.55/0.35–0.40; II 1.05–1.28/0.85–0.98 + 0.50–0.63/0.48–0.50 + 0.78–0.85/0.53–0.63 + 0.70–0.8 8/0.50–0.60 + 0.45/0.35–0.38; III 1.13–1.38/0.98–1.15 + 0.45–0.63/0.45–0.50 + 0.80–0.98/0.65–0.75 + 0.95–1.1 3/0.78–0.90 + 0.43–0.50/0.43–0.45; IV 1.25–1.55/1.03–1.25 + 0.48–0.55/0.35–0.53 + 0.95–1.15/0.65–1.00 + 1.10–1.3 3/0.73–1.10 + 0.35–0.43/0.35–0.55. — Male. Carapace yellow, eye surrounded by black. Translucent scales in vicinity of eyes. On sides of carapace wide band composed of brownish scales. Sternum yellow centrally, brown marginally. Labium yellow with brown basal part. Chelicerae yellow with brownish oblique lines on front sides. Abdomen yellow with two broad longitudinal stripes dorsally, the same stripes on sides, ventrally single line. Spinnerets yellow. Legs yellow, all femora with brown longitudinal stripes. Tibial apophysis of pedipalp straight, embolus short, curved (Fig. 32). — Female. Colouration as in male, but a little paler. Legs yellow. Epigyne with dichotomous pocket in vicinity of epigastric furrow (Fig. 33A–B). Insemination ducts straight, rather long, spermathecae spherical (Fig. 33C).

Habitat. Collected on grass in pine woods.

Distribution. Eastern Palearctic species, new for Khabarovskii Krai. The species was described from Korea (Wesołowska 1981a) on the basis of the female. In the Russian Far East recorded from Primore (the females only) by Prószyński (1979: figs. 154–155 — as "*Icius sp.*"), Dunin




Fig. 32. *Phintella parva* (Wesołowska), male, palpal organ. — A: ventral view. — B: lateral view.

(1984) and Nenilin (1985). The male has been mentioned as *Phintella difficilis* (Bösenberg et Strand) from Primore (Prószyński 1979: figs. 145–149, Dunin 1984, Nenilin 1985) and Furugelm island (Shternbergs 1988).

P. parva is new for Japan; by Bohdanowicz & Prószyński 1987 erroneously reported as *Phintella mellotei* (Simon).

Phintella popovi (Prószyński, 1979)

Fig. 34

Material: Khabarovskii Krai, BR, 2♂, 3♀, 16–18.VI.1987, D. Logunov (MNH), 1♀ (BI, No 772); environs of Khabarovsk, 1♂, 19.VI.1931, V. Sychevskaya (ZMMU); Evreiskaya Autonoma Oblast, Pashkovo village, 3♀, 24.VI.1978, S. Toms (ZI); Nanaiskii distr., Slavvanka, 1♂, 1♀, 28.V.1989, D. Kurenshchikov (BI, No 775); the Amur river, Slavyanskii island, 1♂, 23.VI.1989, D. Kurenshchikov (BI, No 776); Amurskaya Oblast, Khinganskii reserve, Antonovskoe forest-range, 4♀, 2–3.VIII.1985, Yu. Marusik (ZI); Amurskii distr., Nizhnepokrovskoe village, 1♂ (palp only), 25.VI.1988, D. Kurenshchikov (BI, No 777); Primore, Prikhankaiskii distr.,




Fig. 33. *Phintella parva* (Wesołowska), female. — A–B: epigyne. — C: internal structures of epigyne.

Khorol village, 1♂, 18.VI.1981 (ZMMU); Pozharskii distr., middle flow of the Bikin river, 1♂, VIII.1977, Shibnev (BI, No 773); Khazanskii distr., Kedrovaya Pad reserve, 3♀, 16.V–20.VI. 1968, F. Popov (ZI); Vladivostok, Orlineoe Gnezdo, 1♀, IV–V.1903, N. Palchevskii (ZI); Partizanskii distr., Tigrovaya village, 1♀, 12.VI.1927, Martynov (ZI, n — 554-927); Spasskii distr., Yakovlevka village, 1♀, VI.1926, A. Dyakonov, N. Filipov (ZI, n — 539-926); Ussuri area, Kamenushka, 1♀, 7.VI.1981, G. Belova (ZI); Furugelm island, 1♀, 18.VII.1975, M. Shternbergs (BI, No 774); South of Krasnoyarskii Krai, Khakassya, Minnusinsk, 2♀, 1981, Roschier (ZMT).

Diagnosis. Similar to *Phintella parva* (Wesołowska), but may be distinguished by colouration: *P. popovi* has four and *P. parva* two stripes on abdomen. The male easily distinguished by shape of tibial apophysis and embolus (cf. Figs. 34A–B and 32), the female by shorter insemination ducts (cf. Figs. 34D and 33C). The male is easy to recognize, but the female is often confused with other *Phintella* species. Thus, females of *P. popovi* were recorded as *P. linea* from Korea (Wesołowska 1981a) and the Russian Far

East (Nenilin 1985), and also as *Phintella abnormis* (Bösenberg et Strand) from the Russian Far East (Prószynski 1976: map 99, 1979, Dunin 1984: fig. 27).

Description. Measurements (male/female). Cephalothorax: length 1.70–1.88/1.70–2.03, width 1.20–1.30/1.28–1.43, height 0.78/0.75–0.83. Abdomen: length 1.85–1.93/3.13–3.55, width 0.95–1.05/2.10–2.25. Eye field: length 0.80–0.93/0.90–0.93, AL 1.03–1.10/1.08–1.18, PL 1.00–1.08/1.03–1.13, AM 0.33/0.38. Legs: I 1.18–1.53/1.00–1.10+0.55–0.68/0.48–0.55+1.00–1.13/0.73–0.78+0.73–0.85/0.60–0.70+0.48–0.50/0.38–0.45; II 1.03–1.13/0.98–1.05+0.50–0.55/0.48–0.58+0.73–0.83/0.63–0.70+0.83/0.55/0.68+0.40–0.45/0.35–0.38; III 1.10–1.18/1.03–1.25+0.48–0.50/0.40–0.50+0.80–0.88/0.70–0.73+0.95–1.05/0.85–1.05+0.48–0.50/0.43–0.45; IV 1.25–1.38/1.25–1.43+0.43–0.68/0.48–0.58+0.95–1.08/0.90–1.05+1.15–1.45/1.08–1.43+0.45–0.50/0.45–0.50. — Male. Carapace yellow with black margins, eyes surrounded by black. In vicinity of eyes translucent




Fig. 34. *Phintella popovi* (Prószyński), male and female. — A–B: palpal organ, ventral and lateral views. — C–D: epigyne and its internal structures.

scales. Sternum, labium, maxillae and chelicerae yellow. Abdomen yellow with two pairs of longitudinal blackish stripes. Spinnerets yellow. Legs yellow, femora I–III with black spots (see also Prószyński 1979: fig. 150). Pedipalp yellow, its structure shown in Fig. 34A–B. — Female. Colouration resembles male, but a little lighter. Legs yellow. Epigyne and its internal structures given in Fig. 34C–D.

Habitat. Collected from shrubs and grass.

Distribution. Eastern Palaearctic species, new for Khabarovskii Krai, distributed westwards to the Jenisej river.

Most likely all the existing records of *P. abnormis* from the Russian Far East (perhaps with exception of males shown in Dunin 1984: figs. 25–26) refer in fact to *P. popovi*.

***Pseudicius vulpes* (Grube, 1861)**

Material: Khabarovskii Krai, BR, 2♀, 5.VI.1987, D. Logunov (BI); Nanaiskii distr., the Amur river, Slavyanskii island, 1♀, 20.VI.1989, D. Kurenschchikov (MNH).

Habitat. Collected in deciduous woods (aspen, birch, oak) and on meadows.

Distribution. Eastern Palaearctic species, previously recorded from the Russian Far East by Grube (1861), Prószyński (1971, 1976: map 111, 1979), Dunin (1984) and Nenilin (1985).

***Salticus cingulatus* (Panzer, 1797)**

Material: Khabarovskii Krai, Selemdzhinsk, 2♀, VI–VII.1985, A. Kudryavtsev (ZMMU); Amurskaya Oblast, Zeiskii reserve, 1♂, 23.VI.1978, K. Eskov (ZMMU).

Distribution. New for Khabarovskii Krai. Species known from Europe and Western Siberia; recorded from the Russian Far East by Nenilin (1985).

***Sitticus cutleri* Prószyński, 1980**

Material: Khabarovskii Krai, BR, 1♀, 19.VI.1987, D. Logunov (MNH); Okhotskii distr., the Tirbikan stream valley, basin of the Ulya river, 1♀, 20.VIII–14.IX.1986, I. Sukacheva (ZMMU); South of Krasnoyarskii Krai, Khakassya, Shirkii distr., 3–5 km E of Shira village, Itkul lake, 4♂, 21–22.VII.1990, D. Logunov (BI, No 412).

Habitat. Collected in moist glades.

Distribution. New for Khabarovskii Krai. Rare species. Described from Nearctic region (Prószyński 1980). In Russia hitherto recorded from Evenkiya (Western Siberia) (Nenilin 1984, 1985 and Eskov 1988) and Magadanskaya Oblast (Marusik 1988).

Sitticus fasciger (Simon, 1880)

Material: Khabarovskii Krai, BR, Bychikha village, 1♂, 1♀, VII–VIII.1987, S. Ivanov (BI, No 899); 1♀, V.1989 (MNH); Lasovskii distr., shore of the Kiya river, 1♂, VII.1987, Dyachenko (IWP).

Distribution. New for Khabarovskii Krai. In the Russian Far East recorded from Primore (Prószyński 1976: map 172, 1979).

Sitticus floricola (C. L. Koch, 1837)

Material: Khabarovskii Krai, BR, 7♀, 16–17.VI.1987, D. Logunov (BI, No 897), 2♀ (MNH).

Habitat. Collected by sweeping on meadows and in moist places with *Stratiotes* sp. and sparse trees.

Distribution. Widely distributed Holarctic species. Recorded from the Russian Far East by Prószyński (1976: map 190, 1979), Dunin (1984) and Nenilin (1985).

Synageles venator (Lucas, 1836)

Material: Khabarovskii Krai, Nikolaevsk na Amure, 1♂ (incorrectly determined as "*Myrmarachne formicaria*"), [leg.] Schrenck, coll. Grube (MNH); Primore, Khasanskii distr., upper flow of the Kedrovka river, 1♂, 24.VIII.1968, F. Popov (ZI); Ussuriiskii distr., Suptinskii reserve, 1♀, 9.VII.1937, A. Rikhter (ZI, n — 292-1937).

Distribution. Widely distributed Palaearctic species, recorded from the Russian Far East by Prószyński (1976: map 204, 1979 — on basis of the same specimens).

Yaginumaella medvedevi Prószyński, 1979

Material: Khabarovskii Krai, BR, 1♂, 20♀, 6–19.VI.1987, D. Logunov (BI, No 709, 710, 717), 3♂, 5♀ (MNH); 1♀, 2.VI.1988, S. Ivanov (BI, No 716); Khabarovsk re-

gion, Khekhtsir ridge, 1♂, 1♀, VI.1990, D. Kurenschikov (IWP); Primore, Ussuriiskii reserve, 1♂, 1♀, 25.VI.1977, G. Kurcheva (ZMMU); Kedrovaya Pad reserve, 2♂, 29.V–21.VIII.1976, B. Zakharov (BI, No 708, 742), 1♂, 1♀, 25.VII–16.IX.1978, B. Zakharov (BI, No 714, 718).

Habitat. Collected by sweeping in various woods: in deciduous forest (aspen, birch, oak), in deciduous woods with larch, as well as in pine woods. Also found on moister places with single alders.

Distribution. Eastern Palaearctic species, recorded from the Russian Far East from Khabarovskii Krai, Primore, Sakhalin and Kunashir islands (Prószyński 1979, Dunin 1984 and Nenilin 1985).

Yaginumaella striatipes (Grube, 1861)

Fig. 35

Yaginumaella ususudi (Yaginuma, 1972) syn. n.

Material: Khabarovskii Krai, BR, 23♂, 42♀, 8–13.VI.1987, D. Logunov (BI, No 670, 671, 673, 676, 677), 2♂, 15♀ (MNH); Nanaiskii distr., Slavyanka, 2♀, 29.VI.1990, D. Kurenschikov (IWP); Primore, Chuguevskii distr., 1♀, 23.IX.1974, G. Kurcheva (ZMMU). — Additional material: Japan, Kanayama Yamanashi, 5♂, 3♀, 15.VII.1970, M. Ohno, coll. Yaginuma (IBS).

Habitat. Collected in litter and by sweeping in various woods: in coniferous woods (spruce and fir), in mixed woods (pine, birch, rich undergrowth), as well as in deciduous woods (elm). Also found on clearings. *Y. medvedevi* and *Y. striatipes* differ to some extent in their altitudinal distribution. The former occurs in low positions between 100 and 400 m a.s.l. while the latter is found from 250 to above 900 m a.s.l. (highest points of the Bolshekhkhtsirsky reserve).

Distribution. Eastern Palaearctic species, recorded from the Russian Far East by Grube (1861), Prószyński (1971, 1976: map 146, 1979), Dunin (1984) and Nenilin (1985).

In spite of having a series of *Y. ususudi* specimens, in neither sex could we find any consistent difference from *Y. striatipes*, except for minor difference in the structure of tibial apophysis (cf. Fig. 35C–D and 35A–B). Thus, we conclude that they are conspecific, and that *Y. ususudi* constitutes a junior synonym of *Y. striatipes*.




Fig. 35. *Yaginumaella striatipes* (Grube), male, tibial apophysis. — A–B: specimens from Khabarovskii Krai. — C–D: specimens from Japan.

Acknowledgements. We are very grateful to Dr. Yu. M. Marusik (Institute of the Biological Problems of the North, the Russian Academy of Sciences, Magadan), Mr. D. K. Kurenshchikov (Institute of Water and Ecological Problems, the Russian Academy of Sciences, Khabarovsk), Dr. M. T. Shternbergs (Department of Zoology, Latvian University, Riga, Latvia) and Mr. S. I. Ivanov (Bolshekhokhtsirskii reserve) for the loan of their *Salticidae* collections for our study. We also wish to thank Dr. K. G. Mikhailov (Zoological Museum of Moscow University), Dr. S. Koponen (Zoological Museum of University of Turku, Finland) and Dr. V. I. Ovcharenko (Zoological Institute, the Russian Academy of Sciences, St. Petersburg) for giving us the opportunity to study their museum collections. We are especially grateful to Prof. J. Prószyński (Institute of Biology, WSRP, Siedlce, Poland) for his warm

support and discussion of some problems during the preparation of this paper.

References

- Azheganova, N. S. & Stenchenko, T. I. (Ажеганова, Н. С. & Стенченко, Т. И.) 1977: [On the spider fauna of Amur Oblast.] (In Russian) — In: Azheganova, N. (Ажеганова, Н.) (ed.), [Problems of arachnology. Fauna and ecology of spiders.]: 106–111. Mezvuzovskii sbornik. Perm.
- Bohdanowicz, A. & Prószyński, J. 1987: Systematic studies on East Palaearctic Salticidae (Araneae), IV. Salticidae of Japan. — *Ann. Zool.* 41:43–151.

- Bösenberg, W. & Strand, E. 1906: Japanische Spinnen. — *Abh. Senck. Naturf. Ges.* 30:93–422.
- Chikuni, Y. 1989: Some interesting Japanese spiders of the families Amaurobiidae, Araneidae and Salticidae. — In: Nishikawa, Y. & Ono, H. (eds.), *Arachnol. Pap. presented to T. Yaginuma in the occasion of his retirement, Arachnologists' group*: 133–159. Osaka.
- Chikuni, Y. & Yaginuma, T. 1976: *Marpissa magister* and *Marpissa elongata*. (In Japanese) — *Atypus* 67:33–35.
- Dunin, P. M. (Дунин, П. М.) 1984: [Materials on the spider fauna from the Far East (Arachnida, Araneae). Family Salticidae.] (In Russian) — In: Lev, P. (Лев, П.) (ed.), [Fauna and ecology of insects of Southern part of the Far East.]: 128–140. *Sbornik nauchnykh trudov. Vladivostok*.
- Eskov, K. Yu. (Еськов, К. Ю.) 1988: [Araneae of Central Siberia.] (In Russian) — In: Rogacheva, E. (Рогачева, Е.) (ed.), [Faunistic materials from Middle Siberia and neighbouring regions of Mongolia.]: 101–155. Moscow.
- Grube, A. E. 1861: Beschreibung neuer, von den Herren L. v. Schrenck, Maack, C. v. Dittmar u. a. im Amurlande und in Ostsibirien gesammelter Araneiden. — *Bull. Acad. Imp. Sci. S.Petersburg* 4:161–180.
- Izmailova, M. V. (Измайлова, М. В.) 1980: [Spiders of Charskaya Kotlovina.] (In Russian) — In: Kulik, S. (Кулик, С.) (ed.), [Arthropoda of the Siberia and Far East.]: 108–112. Irkutsk.
- Karsch, F. 1879: Baustoffe zu einer Spinnenfauna von Japan. — *Verh. Naturh. Ver. Preuss. Rheinl. Westf.* 36:57–105.
- Kharitonov, D. E. (Харитонов, Д. Е.) 1932: [Catalogue of spiders of the USSR.] (In Russian) — *Ezhгод. Zool. Mus. AN SSSR*, 32: suppl. 206 pp.
- Kulczyński, W. 1895: Attidae Musei Zoologici Varsoviensis in Siberia orientali collecti. — *Rozpr. Spraw. Wyzd. Mat. Przyr. Akad. Umiej.* 32:1–54.
- 1926: *Arachnoidea Samschdhalica*. — *Ann. Mus. Zool. Acad. Sci. USSR*. 27:29–72.
- Logunov, D. V. (Логунов, Д. В.) 1991: The spider family Salticidae (Aranei) from Touva. I. Six new species of the genera *Sitticus*, *Bianor* and *Dendryphantes*. (In Russian with English summary) — *Zool. Zh.* 70:50–60.
- Logunov, D. V. & Marusik, Yu. M. (Логунов, Д. В. & Марусик, Ю. М.) 1991: Redescriptions and morphological differences of *Bianor aurocinctus* (Ohlert) and *B. aemulus* (Gertsch) (Aranei, Salticidae). (In Russian with English summary) — *Sibir. Biol. Zh.* 2:39–47.
- Marusik, Yu. M. (Марусик, Ю. М.) 1988: New species of spiders (Aranei) from the upper Kolyma. (In Russian with English summary) — *Zool. Zh.* 67:1469–1482.
- Marusik, Yu. M. & Logunov, D. V. (Марусик, Ю. М. & Логунов, Д. В.) (in press): Little known species of spiders of the families Salticidae and Thomisidae (Aranei) from the Far East of the USSR. — *Sibir. Biol. Zh.*
- Matsumoto, S. 1989: Colour variation in the prolateral side of carapace and appendages of jumping spider of the genus *Phintella* (Araneida: Salticidae). — In: Nishikawa, Y. & Ono, H. (eds.), *Arachnol. Pap. presented to T. Yaginuma in the occasion of his retirement, Arachnologists' group*: 123–131. Osaka.
- Nenilin, A. B. (Ненилин, А. Б.) 1984: On the taxonomy of spiders of the family Salticidae of the fauna of the USSR and neighbouring countries. (In Russian with English summary) — *Zool. Zh.* 63:1175–1180.
- 1985: [Materials on the fauna of the spider family Salticidae of the USSR. II. Results of the study in the USSR.] (In Russian) — In: Ovcharenko, V. (Овчаренко, В.) (ed.), [Fauna and ecology of spiders of the USSR.]: 129–134. Leningrad.
- Oligier, T. I. (Олигер, Т. И.) 1984: [Materials on the spiders of Lasovskii reserve.] (In Russian) — In: Utochkin, A. (Уточкин, А.) (ed.), [Fauna and ecology of Arachnids.]: 120–127. Univ. of Perm.
- Paik, K. Y. 1987: Studies on the Korean Salticid (Araneae). III. Some new record species from Korea or South Korea and supplementary describe for two species. — *Korean Arachnol.* 3:3–21.
- Prószyński, J. 1971: Redescriptions of the A. E. Grube's East Siberian species of Salticidae (Aranei) in the collection of the Wrocław Zoological Museum. — *Ann. Zool.* 28:205–226.
- 1973: Systematic studies on East Palaearctic Salticidae. II. Redescriptions of Japanese Salticidae of the Zoological Museum in Berlin. — *Ann. Zool.* 30:97–128.
- 1976: Studium systematyczno-zoogeograficzne nad rodziną Salticidae (Aranei) Regionów Palearktycznego i Nearktycznego. — *Rozpr. WSP*, 6. Siedlce. 260 pp.
- 1978: Distributional patterns of the Palaearctic Salticidae (Araneae). — *Proc. Zool. Soc.* 42:335–343.
- 1979: Systematic studies on East Palaearctic Salticidae. III. Remarks on Salticidae of the USSR. — *Ann. Zool.* 34:299–369.
- 1980: Revision of the spider genus *Sitticus* Simon, 1901 (Aranei, Salticidae) IV. *Sitticus floricola* (C. L. Koch) group. — *Ann. Zool.* 36:1–35.
- 1983: Tracing the history of a genus from its geographical range by the example of *Sitticus* (Arachnida: Araneae: Salticidae). — *Verh. Naturwiss. Ver. Hamburg* 26:161–179.
- Schenkel, E. 1963: *Ostasiatische Spinnen aus dem Muséum d'Histoire Naturelle de Paris*. — *Mem. Mus. Hist. Nat.* 25:289–481.
- Shternbergs, M. T. (Штернбергс, М. Т.) 1988: [Materials on the spider fauna of the Primore.] (In Russian) — In: Utochkin, A. (Уточкин, А.) (ed.), [Fauna and ecology.]: 92–97. Univ. of Perm.
- Wesołowska, W. 1981a: Salticidae (Aranei) from North Korea, China and Mongolia. — *Ann. Zool.* 36:45–83.
- 1981b: Redescriptions of the E. Schenkel's East Asiatic Salticidae (Aranei). — *Ann. Zool.* 36:127–160.
- Žabka, M. 1985: Systematic and zoogeographic study on the family Salticidae (Araneae) from Viet-Nam. — *Ann. Zool.* 39:1–485.